

Listing of the Claims:

1. (Currently amended) A method for producing a plurality of dendritic cell/tumor cell hybrids which induce an anti-tumor response when applied to a patient causing a reduction in of the number of tumor cells in said patient, said method comprising:
 - (a) providing dendritic cells,
wherein said dendritic cells are selected from the group consisting of:
~~dendritic cells which are autologous with respect to said patient~~, dendritic cells which are allogeneic with respect to said patient, and dendritic cells which are both allogeneic with respect to said patient and HLA compatible with said patient, and
wherein said dendritic cells are either isolated from bone marrow, lymph or blood, or are differentiated in vitro from dendritic cell precursors isolated from bone marrow, lymph or blood, and
 - (b) fusing said dendritic cells with tumor cells to produce a plurality of dendritic cell/tumor cell hybrids, wherein said dendritic cell is not a T-lymphocyte or B-lymphocyte, wherein said tumor cell is selected from the group consisting of: an autologous tumor cell with respect to said patient, and an allogeneic tumor cell characteristic of the same cancer type with respect to said patient,
 - (c). selecting from said plurality of dendritic cell/tumor cell hybrids that exhibit dendritic cell markers, tumor associated antigens and the capacity to activate naïve T cells in vitro that can recognize the ~~cancer~~ tumor cells of step (b).
2. (Previously presented) The method of claim 1 wherein the dendritic cells of step (a) are produced by culturing said dendritic cell precursors in the presence of cytokines.
3. (Canceled)
4. (Canceled)
5. (Previously presented) The method of claim 1 wherein the dendritic cell of step (a) is of myeloid origin.
6. (Previously presented) The method of claim 1 wherein the dendritic cell of step (a) is of lymphoid origin.
7. (Previously presented) The method of claim 1 wherein the dendritic cell of step (a) is an isolated dendritic cell.

8. (Canceled)
9. (Previously presented) The method of claim 1 wherein said fusing is carried out using PEG.
10. (Canceled)
11. (Canceled)
12. (Canceled)
13. (Canceled)
14. (Canceled)
15. (Cancelled)
16. (Canceled)
17. (Canceled)
18. (Canceled)
19. (Canceled)
20. (Canceled)
21. (Canceled)
22. (Canceled)
23. (Canceled)
24. (Canceled)
25. (Canceled)
26. (Canceled)
27. (Canceled)
28. (Canceled)
29. (Canceled)
30. (Canceled)
31. (Canceled)
32. (Canceled)
33. (Canceled)
34. (Currently amended) The method of claim 1, wherein said tumor cells ~~of step (a)~~ are sensitive to a drug, said method further comprising, after step (c), killing unfused tumor cells by exposure to said drug.
35. (Original) The method according to claim 34 wherein said drug is hypoxanthine-aminopterin-thymidine (HAT).

36. (Canceled)
37. (Canceled)
38. (Canceled)
39. (Canceled)
40. (Canceled)
41. (Canceled)
42. (Previously presented) A method of claim 1 wherein the plurality of dendritic cell/tumor cell hybrids is further induced to express dendritic cell characteristics.
43. (Canceled)
44. (Canceled)
45. (Canceled)
46. (Original) A method of claim 42 wherein said induction is performed using GM-CSF, IFN, γ , TNF- α or a combination thereof.
47. (Canceled)
48. (Canceled)
49. (Canceled)
50. (Previously presented) A method of claim 1 wherein the plurality of dendritic cell/tumor cell hybrids is treated to prevent further proliferation before using it for the induction of an anti-tumor response.
51. (Canceled)
52. (Canceled)
53. (Canceled)
54. (Original) A method of claim 50 wherein said treatment occurs by irradiation.
55. (Canceled)
56. (Canceled)
57. (Canceled)
58. (Previously presented) The method of claim 1, wherein the dendritic cells are from a dendritic-like cell line.
59. (Previously presented) The method of claim 1, wherein the tumor cells of step (b) are from a tumor cell line having at least one tumor associated antigen in common with said tumor sample.

60. (Previously presented) The method of claim 1, wherein the dendritic cell/tumor cell hybrids are hybridomas.
61. (Previously presented) The method of claim 58, wherein said tumor cells of step (b) are from said patient.
62. (Previously presented) The method of claim 59, wherein the tumor cell line is derived from tumor cells from said patient.
63. (Cancelled)
64. (Previously presented) The method of claim 1, wherein said dendritic cells are dendritic-like cells.
65. (Previously presented) The method of claim 1, wherein said tumor cell is an allogeneic tumor cell with respect to said patient, and has one or more tumor associated antigens in common with that of said autologous tumor cell.
66. (Previously presented) the method of claim 1, wherein said anti-tumor response comprises the in vivo induction of immune effectors that confer resistance to a subsequent challenge with tumor cells.